



All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books
Search PubMed for Go Clear

Limits Preview/Index History Clipboard Details

About Entrez

Display Abstract Show 20 Sort by Send to

Text Version

All: 1 Review: 0

Entrez PubMed
Overview
Help | FAQ
Tutorial
New/Noteworthy
E-Utilities

1: J Neurochem. 2004 May;89(3):581-92.

Related Articles, Links



The nerve growth factor precursor proNGF exhibits neurotrophic activity but is less active than mature nerve growth factor.

Fahnestock M, Yu G, Michalski B, Mathew S, Colquhoun A, Ross GM, Coughlin MD.

PubMed Services
Journals Database
MeSH Database
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
Special Queries
LinkOut
My NCBI (Cubby)

Department of Psychiatry and Behavioural Neurosciences, McMaster University, Hamilton, Ontario, Canada. fahnest@mcmaster.ca

Related Resources
Order Documents
NLM Mobile
NLM Catalog
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

Nerve growth factor (NGF) promotes neuronal survival and differentiation and stimulates neurite outgrowth. NGF is synthesized as a precursor, proNGF, which undergoes post-translational processing to generate mature beta-NGF. It has been assumed that, in vivo, NGF is largely processed into the mature form and that mature NGF accounts for the biological activity. However, we recently showed that proNGF is abundant in CNS tissues whereas mature NGF is undetectable, suggesting that proNGF has biological functions beyond its role as a precursor. To determine whether proNGF exhibits biological activity, we mutagenized the precursor-processing site and expressed unprocessed, cleavage-resistant proNGF protein in insect cells. Survival and neurite outgrowth assays on murine superior cervical ganglion neurons and PC12 cells indicated that proNGF exhibits neurotrophic activity similar to mature 2.5S NGF, but is approximately fivefold less active. ProNGF binds to the high-affinity receptor, TrkA, as determined by cross-linking to PC12 cells, and is also slightly less active than mature NGF in promoting phosphorylation of TrkA and its downstream signaling effectors, Erk1/2, in PC12 and NIH3T3-TrkA cells. These data, coupled with our previous report that proNGF is the major form of NGF in the CNS, suggest that proNGF could be responsible for much of the biological activity normally attributed to mature NGF in vivo.

PMID: 15086515 [PubMed - indexed for MEDLINE]

Display Abstract Show 20 Sort by Send to

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

BEST AVAILABLE COPY